



**GCE AS/A LEVEL**

**2300U10-1**

**WEDNESDAY, 17 MAY 2023 – MORNING**

**MATHEMATICS – AS unit 1**

**PURE MATHEMATICS A**

**2 hours 30 minutes plus your additional  
time allowance**

## **ADDITIONAL MATERIALS**

**In addition to this examination paper, you will need:**

- **a WJEC pink 16-page answer booklet;**
- **a Formula Booklet;**
- **a calculator.**

## **INSTRUCTIONS TO CANDIDATES**

**Use black ink, black ball-point pen or your usual method.**

**Answer ALL questions.**

**Write your answers in the separate answer booklet provided, following the instructions on the front of the answer booklet.**

**(Turn over)**

**Use both sides of the paper. Please only write within the white areas of the booklet.**

**Write the question number in the left-hand margin at the start of each answer. Write the sub parts, e.g. (a), (b) and (c), within the white areas of the booklet.**

**Leave at least two line spaces between each answer.**

**Sufficient working must be shown to demonstrate the MATHEMATICAL method employed.**

**Answers without working may not gain full credit.**

**Unless the degree of accuracy is stated in the question, answers should be rounded appropriately.**

**(Turn over)**

**INFORMATION FOR CANDIDATES**

**The maximum mark for this paper is 120.**

**The number of marks is given in brackets at the end of each question or part-question.**

**You are reminded of the necessity for good English and orderly presentation in your answers.**

**(Turn over)**

**ADDITIONAL FORMULAE FOR 2023****LAWS OF LOGARITHMS**

$$\log_a x + \log_a y \equiv \log_a (xy)$$

$$\log_a x - \log_a y \equiv \log_a \left( \frac{x}{y} \right)$$

$$k \log_a x \equiv \log_a (x^k)$$

**(Turn over)**

**SEQUENCES**

**General term of an arithmetic progression:**

$$u_n = a + (n - 1) d$$

**General term of a geometric progression:**

$$u_n = ar^{n-1}$$

**(Turn over)**

**MENSURATION**

**For a circle of radius,  $r$ , where an angle at the centre of  $\theta$  radians subtends an arc of length  $S$  and encloses an associated sector of area  $A$  :**

$$s = r\theta$$

$$A = \frac{1}{2} r^2 \theta$$

# CALCULUS AND DIFFERENTIAL EQUATIONS

## DIFFERENTIATION

**FUNCTION**

**DERIVATIVE**

**$f(x)g(x)$**

**$f'(x)g(x) + f(x)g'(x)$**

**$f(g(x))$**

**$f'(g(x))g'(x)$**

## INTEGRATION

**FUNCTION**

**INTEGRAL**

**$f'(g(x))g'(x)$**

**$f(g(x)) + c$**

Area under a curve =  $\int_a^b y dx$

**(Turn over)**

**REMINDER: Sufficient working must be shown to demonstrate the MATHEMATICAL method employed.**

**01 (a)**

**Using the binomial theorem, write down and simplify the first three terms in the expansion of  $(1 - 3x)^9$  in ascending powers of  $x$  [3 marks]**

**(b) Hence, by writing  $x = 0.001$  in your expansion in part (a), find an approximate value for  $(0.997)^9$  Show all your working and give your answer correct to three decimal places. [3 marks]**

**(Turn over)**

- 02. Solve the following equation for values of  $\theta$  between  $0^\circ$  and  $360^\circ$**   
**[7 marks]**

$$3 \sin^2 \theta - 5 \cos^2 \theta = 2 \cos \theta$$

**03.** The point ***A*** has coordinates  **$(-2, 5)$**  and the point ***B*** has coordinates  **$(3, 8)$** . The point ***C*** lies on the ***X***-axis such that ***AC*** is perpendicular to ***AB***

**(a)** Find the equation of ***AB***

[3 marks]

**(b)** Show that ***C*** has coordinates  **$(1, 0)$**

[3 marks]

**(c)** Calculate the area of triangle ***ABC***

[4 marks]

**(d)** Find the equation of the circle which passes through the points ***A***, ***B*** and ***C***

[5 marks]

(Turn over)

**04 (a)**

Find the remainder when the polynomial  $3x^3 + 2x^2 + x - 1$  is divided by  $(x - 3)$

**[3 marks]**

**(b)** The polynomial  $f(x) = 2x^3 - 3x^2 + ax + 6$  is divisible by  $(x + 2)$ , where  $a$  is a real constant.

**(i)** Find the value of  $a$  **[3 marks]**

**(ii)** Showing all your working, solve the equation  $f(x) = 0$  **[4 marks]**


**(Turn over)**

05. Simplify the expression

$$\sqrt[3]{512a^2} - \frac{a^{\frac{7}{6}} \times a^{-\frac{1}{3}}}{a^{\frac{1}{6}}}$$

[4 marks]

(Turn over)



06. The diagram opposite shows a triangle  $ABC$

Given that

$$AB = 3, BC = 2\sqrt{5}, AC = 4 + \sqrt{3}$$

find the value of  $\cos \hat{A}BC$

Show all your working and give your answer in the form

$$\frac{(a - b\sqrt{3})}{6\sqrt{5}} \text{ where } a, b \text{ are integers.}$$

[7 marks]

(Turn over)

07. The curve  $C$  has equation  $y = 2x^2 + 5x - 12$  and the line  $L$  has equation  $y = mx - 14$  where  $m$  is a real constant.

(a) Given that  $L$  is a tangent to  $C$

(i) show that  $m$  satisfies the equation

$$m^2 - 10m + 9 = 0$$

[5 marks]

(ii) find the coordinates of the two possible points of contact of  $C$  and  $L$

[6 marks]

(Turn over)

07 (b)

Given instead that  $L$  intersects  $C$  at two distinct points, find the range of values of  $m$

[2 marks]

08. Show, by counter example, that the following statement is false.

“For all positive integer values of  $n$ ,  $n^2 + 1$  is a prime number.”

[3 marks]

(Turn over)

09 (a)

Given that  $y = x^2 - 3x$  find  $\frac{dy}{dx}$

from first principles.

[5 marks]

(b) The function  $f$  is defined by

$$f(x) = 4x^{\frac{3}{2}} + \frac{6}{\sqrt{x}} \quad \text{for } x > 0$$

(i) Find  $f'(x)$

[2 marks]

(Turn over)

**09 (b)(ii)**

**When  $x > k$ ,  $f(x)$  is an increasing function. Determine the least possible value of  $k$**

**Give your answer correct to two decimal places.**

**[4 marks]**

**(Turn over)**

**10. Solve the following equations for values of  $X$**

**(a)  $\ln(2x + 5) = 3$**

**[2 marks]**

**(b)  $5^{2x+1} = 14$**

**[3 marks]**

**(c)**

**$3\log_7(2x) - \log_7(8x^2) + \log_7x = \log_381$**

**[6 marks]**

**(Turn over)**

11. The function  $f$  is defined by

$$f(x) = \frac{8}{x^2}$$

(a) Sketch the graph of  $y = f(x)$

[2 marks]

(b) On a separate set of axes, sketch the graph of  $y = f(x - 2)$

Indicate the vertical asymptote and the point where the curve crosses the  $y$ -axis.

[3 marks]

(Turn over)

11 (c)

Sketch the graphs of  $y = \frac{8}{x}$  and

$$y = \frac{8}{(x-2)^2} \text{ on the same set}$$

of axes.

Hence state the number of roots of the equation

$$\frac{8}{(x-2)^2} = \frac{8}{x}$$

[2 marks]

(Turn over)

12. The position vectors of the points **A** and **B** relative to a fixed origin **O** are given by

$$\underline{\mathbf{a}} = -3\underline{\mathbf{i}} + 4\underline{\mathbf{j}} \quad \underline{\mathbf{b}} = 5\underline{\mathbf{i}} + 8\underline{\mathbf{j}}$$

respectively.

- (a) Find the vector **AB**

[2 marks]

- (b) (i)

Find a unit vector in the direction of **a**

[2 marks]

(Turn over)

12 (b)(ii)

The point **C** is such that the vector **OC** is in the direction of **a**


Given that the length of **OC** is **7** units, write down the position vector of **C**

[1 mark]

(c) Calculate the angle **AOB**

[3 marks]

(Turn over)



13 (a)

Find  $\int \left( 4x^{-\frac{2}{3}} + 5x^3 + 7 \right) dx$

(b) The diagram opposite shows the graph of

$$y = x(x + 6)(x - 3)$$

Calculate the total area of the regions enclosed by the graph and the  $X$ -axis.

[9 marks]

(Turn over)

14 (a)

Two variables,  $X$  and  $y$  are such that the rate of change of  $y$  with respect to  $X$  is proportional to  $y$

State a model which may be appropriate for  $y$  in terms of  $X$

[1 mark]

(b) The concentration,  $Y$  units, of a certain drug in a patient's body decreases exponentially with respect to time. At time  $t$  hours the concentration can be modelled by  $Y = Ae^{-kt}$  where  $A$  and  $k$  are constants.

(Turn over)

**14 (b)**

**A patient was given a dose of the drug that resulted in an initial concentration of 5 units.**

- (i) After 4 hours, the concentration had dropped to 1.25 units. Show that  $k = 0.3466$  correct to four decimal places.**

**[2 marks]**

- (ii) The minimum effective concentration of the drug is 0.6 units. How much longer would it take for the drug concentration to drop to the minimum effective level?**

**[3 marks]**

**END OF PAPER**