



GCE AS/A LEVEL – NEW

2420U20-1

PHYSICS – AS unit 2

Electricity and Light

THURSDAY, 8 JUNE 2017 – AFTERNOON

1 hour 30 minutes plus your additional time allowance

Surname _____

Other Names _____

Centre Number _____

Candidate Number 2 _____

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	6	
2.	8	
3.	8	
4.	11	
5.	12	
6.	10	
7.	12	
8.	13	
Total	80	

ADDITIONAL MATERIALS

In addition to this examination paper, you will require a calculator and a DATA BOOKLET.

INSTRUCTIONS TO CANDIDATES

Use black ink, black ball-point pen or your usual method.

Write your name, centre number and candidate number in the spaces provided on the front cover.

Answer ALL questions.

Write your answers in the spaces provided in this booklet. If you run out of space use the continuation page(s) at the back of the booklet taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES


The total number of marks available for this paper is 80.

The number of marks is given in brackets at the end of each question or part-question.

The assessment of the quality of extended response (QER) will take place in question 5(a).

Answer ALL questions.

1. A strip of metal of rectangular cross-section is shown.



- (a) Calculate the drift velocity of free electrons in the strip when there is a current of 8.0 A as shown. The free electron concentration, n , in the metal is $8.5 \times 10^{28} \text{ m}^{-3}$. [3]

1(b) The resistance of the strip is $0.15\ \Omega$ from end to end. Calculate the energy transferred from electrical potential to thermal in the strip in a time of 20 s, for a current of 8.0 A, AND briefly state how the transfer takes place. [3]


6

2. A student is given wire cutters and a reel of metal wire, and asked to find the resistivity of the metal. She cuts off a piece of the wire and makes measurements on it. She takes repeat readings and obtains the same values each time.

Quantity	Measurement	Instrument	Resolution of instrument
length	812 mm	metre ruler	1 mm
diameter	0.48 mm	digital calipers	0.01 mm
resistance	2.2 Ω	digital multimeter	0.1 Ω

- (a) (i) Calculate the resistivity of the metal of the wire. [3]

2(b) Suggest ONE way in which the student could reduce the uncertainty in her value for the resistivity, using the same reel of wire and the same instruments as before. Explain briefly why the uncertainty would be reduced. [2]



3. For the potential divider circuit shown opposite a student is asked to select values for R_1 and R_2 so that the circuit meets certain requirements.


REQUIREMENTS

- A** an output pd, V_{out} , of 3.0 V when unloaded (switch open),
- B** a power dissipation of less than 0.50 W in R_1 when unloaded (switch open),
- C** a DECREASE of less than 0.40 V in V_{out} when the switch is closed.

Showing your working clearly, determine whether or not the circuit meets the requirements when $R_1 = 450 \Omega$ and $R_2 = 150 \Omega$.

- (a) Requirement **A**: an output pd, V_{out} , of 3.0 V when unloaded (switch open). [2]

3(b) Requirement B: a power dissipation of less than 0.50 W in R_1 when unloaded (switch open). [2]



4(a) (ii) Carefully sketch **ON THE DIAGRAM OPPOSITE** page 13 the same part of the string at time $t = 0.34$ s. (The periodic time is 0.040 s.) [1]

(b) If the string above is clamped rigidly a little way to the right of the part shown, a **STATIONARY** wave may be observed.

(i) State what is meant by a **NODE** in a stationary wave, and state how far apart the nodes will be in **THIS** stationary wave. [2]

4(b) (ii) State how, if at all, the PHASE of the oscillations of the particles of the string varies with distance along the string, for a stationary wave. [2]

4(b) (iii) Explain, in terms of INTERFERENCE, how the stationary wave is produced for this stationary wave on the string. [2]

4(b) (iv) Stationary waves may also be observed on the string when the wave source is set to higher frequencies. State ONE way, apart from having a different frequency, in which the stationary waves will be different from the original stationary waves. [1]

- 5(a) Describe in detail how you would determine the wavelength of light from a (low-power) laser, given an opaque plate with two parallel slits. (The distance between the centres of the slits is less than 1 mm and has already been measured.)**

[6 QER]




DIAGRAM NOT TO SCALE

5(b) A diffraction grating is used with a laser as shown opposite, and a line of bright spots is seen on the screen as shown opposite.


- (i) Using a labelled sketch of a relevant triangle (in the space below), explain why, to a very good approximation:**

$$\sin\theta_1 = \frac{0.0153}{0.600}$$

in which θ_1 is the angle between the zeroth order and first order beams. [2]


5(b) (ii) Determine the wavelength of light from the laser. [2]

- 6(a) A student sets up a simple periscope consisting of two glass prisms ($n_{\text{glass}} = 1.52$). She tests it by using it to view a light emitting diode (LED) as shown.



- (i) Give the full name of the process by which the light changes direction in the prisms.


[1]



7(a) A student uses the apparatus shown in the diagram opposite to determine the maximum kinetic energy, $E_{k\max}$, of electrons ejected from a strontium surface by light of wavelength 405 nm.

He records that $E_{k\max} = 0.480 \text{ eV}$.

(i) Describe briefly how he used the apparatus to obtain this result. [2]



8(a) A simplified energy level diagram for a 3-level laser system is given.

level P _____ $2.03 \times 10^{-19} \text{ J}$

level U _____ $1.28 \times 10^{-19} \text{ J}$

level L _____ 0
(ground state)

- (i) Calculate the wavelength of photons which can take part in transitions between levels U and L. [2]

8(a) (ii) Describe the processes of **ABSORPTION** and **STIMULATED EMISSION** involving levels U and L. [4]

ABSORPTION:

STIMULATED EMISSION:

8(a) (iii) If the relative populations of levels P, U and L are 0.2%, 54.0% and 45.8%, state with a reason whether absorption or stimulated emission is the more likely event for a photon of the wavelength calculated in (a)(i). [1]

8(a) (iv) The laser is pumped by promoting electrons from the ground state to level P. Explain why electrons must spend only a very short time in level P, but a relatively long time in level U. [2]

13

END OF PAPER

