

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

* 7 2 3 4 3 3 2 2 6 *

CHEMISTRY

0620/32

Paper 3 (Extended)

October/November 2015

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use an HB pencil for any diagrams or graphs.

Do not use staples, paper clips, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used.

A copy of the Periodic Table is printed on page 12.

You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.


The syllabus is approved for use in England, Wales and Northern Ireland as a Cambridge International Level 1/Level 2 Certificate.

This document consists of **12** printed pages.


1 Use your copy of the Periodic Table to help you answer some of these questions.

(a) Predict the formulae of the following compounds.

(i) nitrogen fluoride NF_3



(ii) phosphorus sulfide P_2S_3



[2]

(b) Deduce the formulae of the following ions.

(i) selenide Se^{2-} *Se is in grp 6 of the periodic table .So it will have 6 valence electrons. So it will accept 2 electrons when forming an ion*

(ii) gallium Ga^{3+} *Ga is in the group 3 of the periodic table. So it will want to lose 3 electrons when forming an ion*

[2]

(c) Use the following ions to determine the formulae of the compounds.

ions OH^- Cr^{3+} Ba^{2+} SO_4^{2-}



compounds

(i) chromium(III) sulfate $\text{Cr}_2(\text{SO}_4)_3$



(ii) barium hydroxide $\text{Ba}(\text{OH})_2$

[2]

[Total: 6]

2 (a) Polluted air contains two oxides of carbon and two oxides of nitrogen. A major source of these pollutants is motor vehicles.

(i) Describe how carbon dioxide and carbon monoxide are formed in motor vehicle engines.

When fuel in a vehicle burns, then it produces CO_2 in case of complete combustion and

CO in case of incomplete combustion. In this way CO_2 and CO are formed in motor

vehicle engines

[3]

(ii) State one adverse effect of each of these gases.

CO_2 causes global warming

CO is a poisonous gas and may result in health poisoning like tissue damage or worse cases death

[2]

(iii) Nitrogen monoxide, NO , is released by motor vehicle exhausts.

Explain how nitrogen monoxide is formed in motor vehicle engines.

Nitrogen and oxygen combine at high temperature in the car engine to produce NO

[2]

(iv) When nitrogen monoxide is released into the atmosphere, nitrogen dioxide, NO_2 , is formed.

Suggest an explanation why this happens.

NO combines with more oxygen in the atmosphere to form NO_2

[1]

(b) Predict the possible adverse effect on the environment when this non-metal oxide, NO_2 , reacts with water and oxygen.

1. It lowers the pH of the rivers and lakes and kills the aquatic animals.

2. It causes acid rain

[2]

(c) How are the amounts of carbon monoxide and nitrogen monoxide emitted by modern motor vehicles reduced? Include an equation in your answer.

The amount of carbon monoxide and nitrogen oxide emitted by vehicles is reduced by

fitting catalytic converters.

Equation: $2\text{NO} + 2\text{CO} \rightarrow 2\text{CO}_2 + \text{N}_2$

[3]

[Total: 13]

- 3 Two of the main uses of zinc are for galvanising and for making alloys.

One of the main ores of zinc is zinc blende, ZnS. There are two stages in the extraction of zinc from this ore.

- (a) **Stage 1** Zinc oxide is made from zinc blende.

Describe how this is done and write a word equation for the reaction.

Method: zinc blende is roasted heated in air. In this way zinc oxide is formed from zinc blende

Equatio: zinc sulfide + oxygen → zinc oxide + sulfur dioxide;

[2]

- (b) **Stage 2** Zinc oxide is reduced to zinc.

Write a word equation for the reduction of zinc oxide by coke.

zinc oxide + carbon → zinc + carbon dioxide / monoxide;

[1]

- (c) The zinc produced by this process is impure. It can be purified by electrolysis using a method which is similar to the purification of copper. Under the conditions used in the process, zinc is the product at the negative electrode (cathode).

Complete the following description of this purification.

The electrolyte is aqueous zinc sulfate [1]

The negative electrode (cathode) is made of pure zinc [1]

The positive electrode (anode) is impure zinc.

The equation for the reaction at the cathode is $Zn^{2+} + 2e^{-} \rightarrow Zn$; [1]

The equation for the reaction at the anode is $Zn \rightarrow Zn^{2+} + 2e^{-}$; [1]

Explain why the concentration of the electrolyte does not change.

Zinc ions get removed from the solution. At the same time zinc ions are replaced back

into the solution. This happens at the same rate

[2]

(d) Brass is an alloy which contains zinc.

(i) Name the other metal in brass.

Copper

..... [1]

(ii) Suggest two reasons why an alloy such as brass is preferred to either of its constituent metals.

Only 2 points needed

1. It is stronger and harder

2. It has a better appearance. 3. It offers more resistance to corrosion

..... [2]


(e) In an experiment to investigate the rate of rusting of steel, three pieces of steel were used. One piece of steel was completely coated with copper, one piece completely coated with zinc and the third piece was left uncoated. All three pieces were left exposed to the atmosphere.

(i) Explain why the uncoated piece started to rust.

This is because the iron in the steel gets exposed to oxygen and water.

..... [1]

(ii) The coating on both of the other two pieces was scratched, exposing the steel.



The piece of steel coated with zinc still did not rust but the copper-coated piece of steel rusted very rapidly.

Explain these observations in terms of the formation of ions and the transfer of electrons.

Zn more reactive than Fe. Therefore Zn loses more readily and forms (+ve) ions in preference to Fe

Fe is more reactive than Cu; Fe loses electrons more readily and forms (+ve) ions in preference to Cu

..... [4]

[Total: 17]


4 (a) Propane reacts with chlorine to form a mixture of chloropropanes. This is a photochemical reaction.

(i) What is meant by the phrase *photochemical reaction*?


A photochemical reaction is a reaction whose rate is influenced by light or a reaction which occurs in presence of light.

[1]

(ii) The products of this reaction include two isomers, one of which has the following structural formula.



Draw the structural formula of the other isomer.



[1]


(iii) Explain why these two different compounds are isomers.

Both these structures have the same molecular formula but different structural formula

[2]

(b) Bond breaking is an endothermic change and bond forming is an exothermic change.

Bond energy is the amount of energy in kJ/mol needed to break one mole of the specified bond.



Use the following bond energies to determine whether this reaction is exothermic or endothermic. You must show your reasoning.


bond	bond energies in kJ/mol
C-Cl	338
C-H	412
Cl-Cl	242
H-Cl	431
C-C	348

Bond breaking		Bond making	
8 (H-C) bonds are broken =	8 × 412 = 3296 kJ/mol	7 (H-C) bonds formed =	7 × 412 = 3448 kJ/mol
2 (C-C) bonds	= 2 × 348 = 696 kJ/mol	2 (C-C) bonds	= 2 × 348 = 696 kJ/mol
1 (Cl-Cl) bond	= 1 × 242 = 242 kJ/mol	1 (C-Cl) bond	= 1 × 338 = 338 kJ/mol
Total energy supplied	= 4234 kJ/mol	1 (H-Cl) bond	1 × 431 = 431 kJ/mol
		Total energy released	= 4349 kJ/mol

Ans: Energy released is more than energy supplied

- (c) (i) Chloropropane can be hydrolysed to propanol, $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$, by sodium hydroxide.

Write the equation for this reaction.



- (ii) Propanol can be dehydrated. It loses a water molecule to form a hydrocarbon.

Give the name and structural formula of this hydrocarbon.

name propene;

structural formula



[2]

- (iii) Propanol is oxidised to a carboxylic acid by acidified potassium manganate(VII).

Deduce the name of this acid.

propanoic acid

[1]

- (d) Propanol reacts with methanoic acid to form the ester propyl methanoate.



4.0g of methanoic acid was reacted with 6.0g of propanol.

- (i) Calculate the M_r of methanoic acid = 46 [1]
- (ii) Calculate the M_r of propanol = 60 [1]
- (iii) Determine which one is the limiting reagent. Show your reasoning.

moles of $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH} = 0.1$;

moles of $\text{HCOOH} = 0.087$ (0.09)

and limiting reagent is methanoic acid; [2]

- (iv) Calculate the maximum yield in grams of propyl methanoate, $M_r = 88$.

..... $88 \times (\text{mol of limiting reagent in 4(d)(iii)})$; [1]

expected answer: $88 \times 0.087 = 7.65 \text{ g}$;

[Total: 17]

5 Iron is extracted from its ore, hematite, in a blast furnace.

Substances added to the furnace are:

- iron ore, hematite, containing impurities such as silica, SiO_2
- air
- coke, C
- limestone, CaCO_3

Substances formed in the blast furnace are:

- molten iron
- molten slag
- waste gases such as carbon dioxide

(a) State the two functions of the coke used in the blast furnace.

1. It is a source of heat energy

2. It is used as a reducing agent

[2]

(b) Write an equation for the conversion of hematite, Fe_2O_3 , to iron.

(b) $\text{Fe}_2\text{O}_3 + 3\text{CO} \rightarrow 2\text{Fe} + 3\text{CO}_2$ species

[2]

(c) Explain how the silica impurity is removed and separated from the molten iron.

Silica reacts with limestone to form slag. The molten slag forms a layer above the more dense molten iron and they can be both separately, and regularly, drained away.

[3]

(d) The molten iron from the furnace is impure. It contains impurities which include the element carbon.

Explain how the carbon is removed. Include an equation in your answer.

Oxygen is blown over the molten iron. carbon reacts with oxygen and carbon dioxide is formed. carbon dioxide being a gas escapes.

Reaction: $\text{C} + \text{O}_2 \rightarrow \text{CO}_2$

This is how carbon is removed from molten iron

[3]

[Total: 10]

- 6 The table below shows the elements in the third period of the Periodic Table, the number of electrons in their outer energy level, their oxidation state in their common compounds and their melting points.

element	Na	Mg	Al	Si	P	S	Cl	Ar
number of outer electrons	1	2	3	4	5	6	7	8
oxidation state	+1	+2	+3	+4/−4	−3	−2	−1	0
melting point/°C	98	650	660	1414	317	115	−101	−189

- (a) Describe and explain the variation in oxidation state across the period.

The number of e⁻ gained or lost is equal to the numerical value of oxidation state.

- 1) Electrons are lost from Na to Si
- 2) Electrons are gained from Si to Cl
- 3) Si either gains or loses electrons.
- 4) Argon neither gains nor loses electrons

- (b) The first three elements, Na, Mg and Al, are metals.

Describe the structure of a typical metal.

Metals are made of metallic ions which are positively charged. These metallic ions are arranged in a lattice. Also present in the lattice are a sea of electrons, also known as the delocalised electrons.

[3]

- (c) Explain why Na, Mg and Al are good conductors of electricity.

Na, Mg and Al are good conductors because they have free electrons

[1]

- (d) Which element exists as diatomic molecules of the type X₂?

Chlorine

[2]

- (e) Silicon has a similar structure to diamond.

Explain why silicon has the highest melting point in the period.

Silicon is a macromolecule with strong covalent bonds. So it has the highest melting point in the group

[2]

- (f) Sodium chloride is a crystalline solid with a high melting point. It dissolves in water to give a neutral solution. Phosphorus trichloride is a liquid at room temperature. It reacts with water to form an acidic solution.

Suggest an explanation for these differences in properties.

NaCl is an ionic compound and PCl_3 is a covalent compound. NaCl has strong ionic bonds but intermolecular forces are weak. SO it dissolves in water. PCl_3 is a liquid at room temperature because it has weak Vanderwals forces between its molecules.


[3]

- (g) Describe how you could show that magnesium oxide is a basic oxide and not an amphoteric oxide.

Magnesium oxide will neutralise acidic oxide. If MgO is amphoteric then it will also react with a base to neutralise it.

But MgO is not amphoteric, hence it will not react with a base or alkali or basic oxide.

- (h) Draw a dot-and-cross diagram showing the bonding in magnesium oxide. Show outer electrons only.



[3]

[Total: 17]

DATA SHEET
The Periodic Table of the Elements

		Group																																
I	II	III	IV	V	VI	VII	O					O																						
7 Li Lithium 3	9 Be Beryllium 4	1 H Hydrogen 1	12 C Carbon 6	14 N Nitrogen 7	16 O Oxygen 8	19 F Fluorine 9	20 Ne Neon 10	11 B Boron 5	13 Al Aluminium 13	14 Si Silicon 14	15 P Phosphorus 15	17 Cl Chlorine 17	4 He Helium 2																					
23 Na Sodium 11	24 Mg Magnesium 12	26 Fe Iron 26	28 Ni Nickel 28	29 Cu Copper 29	30 Zn Zinc 30	31 Ga Gallium 31	32 Ge Germanium 32	33 As Arsenic 33	34 Se Selenium 34	35 Br Bromine 35	36 Kr Krypton 36	37 Rb Rubidium 37	38 Sr Strontium 38	39 Y Yttrium 39	40 Ca Calcium 20	41 Nb Niobium 41	42 Mo Molybdenum 42	43 Tc Technetium 43	44 Ru Ruthenium 44	45 Rh Rhodium 45	46 Pd Palladium 46	47 Ag Silver 47	48 Cd Cadmium 48	49 In Indium 49	50 Sn Tin 50	51 Sb Antimony 51	52 Te Tellurium 52	53 I Iodine 53	54 Xe Xenon 54					
55 Cs Caesium 55	56 Ba Barium 56	57 La Lanthanum 57	58 Ce Cerium 58	59 Pr Praseodymium 59	60 Nd Neodymium 60	61 Pm Promethium 61	62 Sm Samarium 62	63 Eu Europium 63	64 Gd Gadolinium 64	65 Tb Terbium 65	66 Dy Dysprosium 66	67 Ho Holmium 67	68 Er Erbium 68	69 Tm Thulium 69	70 Yb Ytterbium 70	71 Lu Lutetium 71	72 Hf Hafnium 72	73 Ta Tantalum 73	74 W Tungsten 74	75 Re Rhenium 75	76 Os Osmium 76	77 Ir Iridium 77	78 Pt Platinum 78	79 Au Gold 79	80 Hg Mercury 80	81 Tl Thallium 81	82 Pb Lead 82	83 Bi Bismuth 83	84 Po Polonium 84	85 At Astatine 85	86 Rn Radon 86			
87 Fr Francium 87	88 Ra Radium 88	89 Ac Actinium 89	90 Th Thorium 90	91 Pa Protactinium 91	92 U Uranium 92	93 Np Neptunium 93	94 Pu Plutonium 94	95 Am Americium 95	96 Cm Curium 96	97 Bk Berkelium 97	98 Cf Californium 98	99 Es Einsteinium 99	100 Fm Fermium 100	101 Md Mendelevium 101	102 No Nobelium 102	103 Lr Lawrencium 103	104 Rf Rutherfordium 104	105 Db Dubnium 105	106 Sg Seaborgium 106	107 Bh Bohrium 107	108 Hs Hassium 108	109 Mt Meitnerium 109	110 Ds Darmstadtium 110	111 Rg Roentgenium 111	112 Cn Copernicium 112	113 Nh Nihonium 113	114 Fl Flerovium 114	115 Mc Moscovium 115	116 Lv Livermorium 116	117 Ts Tennessine 117	118 Og Oganesson 118			
133 Cs Caesium 55	137 Ba Barium 56	139 La Lanthanum 57	140 Ce Cerium 58	141 Pr Praseodymium 59	144 Nd Neodymium 60	150 Sm Samarium 62	152 Eu Europium 63	157 Gd Gadolinium 64	159 Tb Terbium 65	162 Dy Dysprosium 66	165 Ho Holmium 67	167 Er Erbium 68	169 Tm Thulium 69	173 Yb Ytterbium 70	175 Lu Lutetium 71	176 Yt Ytterbium 71	177 Hf Hafnium 72	178 Ta Tantalum 73	180 Hg Mercury 80	181 Ta Tantalum 73	182 Hf Hafnium 72	183 W Tungsten 74	184 Re Rhenium 75	186 Os Osmium 76	192 Ir Iridium 77	195 Pt Platinum 78	197 Au Gold 79	201 Hg Mercury 80	204 Tl Thallium 81	207 Pb Lead 82	209 Bi Bismuth 83	210 Po Polonium 84	211 At Astatine 85	212 Rn Radon 86

*58-71 Lanthanoid series
†90-103 Actinoid series

Key

a	X	a = relative atomic mass
	X	X = atomic symbol
b		b = proton (atomic) number

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).