



GCSE

3410UB0-1

MONDAY, 22 MAY 2023 – MORNING

CHEMISTRY – Unit 2:

**Chemical Bonding, Application of Chemical
Reactions and Organic Chemistry
HIGHER TIER**

1 hour 45 minutes plus your additional time allowance

Surname _____

First name(s) _____

Centre Number _____

Candidate Number **0** _____

ADDITIONAL MATERIALS

In addition to this paper you will need a calculator and a ruler.

ITEMS INCLUDED WITH QUESTION PAPER

A separate Data Booklet.

A separate Diagram Booklet.

The Diagram Booklet MUST be handed in to the invigilators and sent for marking.

INSTRUCTIONS TO CANDIDATES

Use black ink, black ball-point pen or your usual method.

Write your name, centre number and candidate number in the spaces provided on the front cover.

Answer ALL questions.

Write your answers in the spaces provided in this booklet. If you run out of space, use the additional pages at the back of the booklet, taking care to number the question(s) correctly.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

Question 6(a) is a quality of extended response (QER) question where your writing skills will be assessed.

The Periodic Table and the formulae for some common ions are printed in the separate Data Booklet.

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
1.	9	
2.	11	
3.	9	
4.	7	
5.	11	
6.	8	
7.	9	
8.	10	
9.	6	
Total	80	

Answer ALL questions

1 (a) A student made some copper(II) sulfate crystals by reacting copper(II) carbonate powder with sulfuric acid using the following method.

Stage 1 Measure 50 cm³ of sulfuric acid into a beaker.

Stage 2 Add copper(II) carbonate powder, one spatula at a time, until all the acid has reacted.

Stage 3 Filter the mixture.

Stage 4 Obtain crystals from the solution.

(i) State how you would carry out Stage 4 to get the largest possible crystals. [1 mark]

1 (a)(ii)

Crystals of copper(II) sulfate could also be made using copper(II) oxide powder instead of copper(II) carbonate powder. State and explain how the observations in Stage 2 would be different. [2 marks]

(iii) Complete the symbol equation for the reaction between copper(II) oxide and sulfuric acid. Copper(II) sulfate is one of the products. [2 marks]



(Turn over)

- 1 (b) DIAGRAM 1.1** in the separate diagram booklet shows an energy profile for a reaction.
- (i)** In **TABLE 1.2** in the separate diagram booklet, give the **LETTER** that represents each part of the energy profile. [2 marks]
- (ii)** Give the meaning of the term activation energy. [1 mark]

- (iii)** State how the energy profile shows that this is an exothermic reaction. [1 mark]

2 (a) Crude oil is separated into mixtures of hydrocarbon compounds in the process of fractional distillation. Many of these fractions are used as fuels. This process is shown in DIAGRAM 2.1 in the separate diagram booklet.

(i) Name the changes of state happening at X and at Y. [1 mark]

X

Y

(ii) Explain why different fractions are formed at different levels. [2 marks]

(Turn over)

2 (a)(iii)

A hydrocarbon fuel was burned and used to heat 100 g of water. The water temperature rose from 18.5 °C to 38.2 °C.

Use the equation below to calculate the amount of energy released by this fuel. Give your answer to TWO significant figures.
[3 marks]

energy (J) = mass of water (g) × 4.2 × temperature rise (°C)

Energy = _____ J

(Turn over)

2 (b) The products of fractional distillation can undergo a process called cracking to produce smaller, more useful hydrocarbons.

(i) Complete the equation for the cracking of $C_{16}H_{34}$. [1 mark]



(ii) State the TWO conditions used for cracking. [1 mark]

(iii) The molecule with the formula C_2H_4 is an unsaturated hydrocarbon.

Give the meaning of the term unsaturated. [1 mark]

(Turn over)

2 (b)(iv)

State why there is a high demand for each of the following products of the cracking reaction. [2 marks]

octane / C_8H_{18}

ethene / C_2H_4

11

3 (a) TABLE 3.1 in the separate diagram booklet shows information about some organic compounds.

(i) COMPLETE THE TABLE. [3 marks]

(ii) The molecular mass of an alkene is 98. Give the molecular formula of this alkene. [1 mark]

$$A_r(\text{C}) = 12 \quad A_r(\text{H}) = 1$$

Molecular formula _____

(Turn over)

- 3 (b) Polyvinyl chloride (PVC) is formed from a monomer, shown in DIAGRAM 3.2 in the separate diagram booklet, in a polymerisation reaction.

Complete the equation for the polymerisation reaction in DIAGRAM 3.3 in the separate diagram booklet. [2 marks]

- (c) A student was asked to draw structures for isomers with the molecular formula C_4H_8 .

She drew the diagrams in DIAGRAM 3.4 in the separate diagram booklet. **Circle** ALL the **CORRECT** structures. [2 marks]

- (d) Pent-2-ene reacts with bromine water in a similar manner to ethene.

The equation for the reaction of pent-2-ene with bromine is shown in DIAGRAM 3.5 in the separate diagram booklet. Complete the equation by drawing the structure of the product. [1 mark]

9

- 4 (a) **TABLE 4.1** in the separate diagram booklet shows the results of an experiment in which zinc and lead powders were added separately to solutions of sodium chloride and iron(II) chloride.

Use the results to place the four metals in order of reactivity. [1 mark]

Most reactive

Least reactive

- (b) A student investigated metal reactivity using a different method.

He added 10.0 g of magnesium powder in 2.0 g portions to 50 cm³ of zinc chloride solution and recorded the temperature of the mixture after each addition. He repeated the experiment with aluminium powder and again with copper powder.

The results for magnesium powder are shown in **TABLE 4.2** in the separate diagram booklet.

question continues on the next page

(Turn over)

4 (b) continued

The results for aluminium and copper have been plotted on **GRAPH 4.3** in the separate diagram booklet.

- (i) Plot the results for magnesium on the grid of **GRAPH 4.3**. Draw a suitable line. [3 marks]
- (ii) State the conclusions that can be drawn about the reactivities of magnesium, aluminium, copper and zinc. Give your reasoning. [3 marks]

7

5 (a)(i) Draw a dot and cross diagram to show the bonding in a carbon dioxide molecule. [2 marks]

carbon (2,4) oxygen (2,6)

(ii) Explain why carbon dioxide has a low boiling point. [2 marks]

continue your answer on the next page

(Turn over)

5 (b)(i) Draw a diagram to show the electronic changes that take place during the formation of magnesium oxide. Include the charges on the ions formed. [2 marks]


magnesium (2,8,2) oxygen (2,6)

5 (b)(ii)

Explain why magnesium oxide has a higher melting point than sodium chloride. [2 marks]

5 (b)(iii)

Sodium chloride is produced by the reaction of sodium with chlorine.



In a reaction using 0.080 mol of sodium, 4.12 g of sodium chloride was produced.

Calculate the percentage yield of this reaction. [3 marks]

$$A_r(\text{Na}) = 23 \quad A_r(\text{Cl}) = 35.5$$

Percentage yield = _____ %

11

(Turn over)

- 6 (b) State how adding sodium hydroxide solution allows solutions containing iron(II) and iron(III) ions to be identified. Give the observations made for both solutions. [2 marks]**

8

7 (a) The formation of ammonia can be represented by the equation in DIAGRAM 7.1 in the separate diagram booklet.

- The total amount of energy released in making the bonds in the products is 2340 kJ
- The total amount of energy released in making the bonds in the products is 94 kJ MORE THAN the total energy used in breaking the bonds in the reactants
- The amount of energy used to break the $\text{N} \equiv \text{N}$ bond is 941 kJ

Use this information to calculate the energy used to break ONE H—H bond. [3 marks]

Energy = _____ kJ

(Turn over)

7 (b) The Haber process used to make ammonia is usually carried out at a temperature of around 400 °C. Explain why this is the optimum temperature. [2 marks]

A higher temperature is not used because

A lower temperature is not used because

(c) Ammonia is used in the production of fertilisers such as ammonium nitrate.

Give the balanced symbol equation for the reaction between nitric acid and ammonia to form ammonium nitrate. [2 marks]

7 (d) Eutrophication is a problem caused by fertilisers being washed into waterways leading to overgrowth of plants. Describe how eutrophication leads to the death of aquatic organisms. [2 marks]

9

- 8 (a) According to the Journal of the British Dental Association, the increase in consumption of fruit juices and other acidic drinks is believed to be one of the leading causes of dental erosion in children and adolescents.

Many fruit juice drinks contain more than one type of acid. Citric acid can be used as a natural preservative and provides a sour taste. Ascorbic acid (vitamin C) is a water-soluble vitamin that must be consumed regularly to ensure proper body function. Citrus fruits, as well as tomatoes and other fresh vegetables, are good sources of vitamin C.

TABLE 8.1 in the separate diagram booklet shows information about the content of different fruit juice drinks.

- (i) Tick (✓) the TWO conclusions that can be drawn from the information. [2 marks]

- as pH increases, citric acid content decreases and sugar content increases
- as acidity decreases, ascorbic acid content decreases and water content decreases
- tomatoes are a good source of vitamin C and citric acid
- citrus fruits contain ascorbic acid and a natural preservative

(Turn over)

8 (a)(ii)

USE INFORMATION FROM TABLE 8.1 to suggest why citric acid content has a much greater effect on the pH than ascorbic acid content. [1 mark]

(b) **The concentration of a solution of sodium hydroxide can be determined by titrating with sulfuric acid. The equation for the reaction is shown.**



(i) **Give the IONIC equation for the formation of water in any neutralisation reaction. Include state symbols. [2 marks]**

(Turn over)

8 (b)(ii)

21.0 cm³ of sulfuric acid with a concentration of 0.350 mol/dm³ neutralised 25.0 cm³ of the sodium hydroxide solution.

- I. Calculate the number of moles of sulfuric acid used in the reaction. [1 mark]

Number of moles = _____ mol

- II. Calculate the concentration of the sodium hydroxide solution. [2 marks]

Concentration = _____ mol/dm³

(Turn over)

8 (b)(iii)

During a similar titration reaction, 0.36 g of water was produced. Calculate the number of MOLECULES of water produced in this reaction.

Give your answer in STANDARD FORM.
[2 marks]

$$A_r(\text{H}) = 1 \quad A_r(\text{O}) = 16$$

$$\text{Avogadro's constant} = 6.0 \times 10^{23}$$

Number of molecules = _____

10

(Turn over)

- 9 Aluminium metal is produced industrially using electrolysis, due to the relatively high stability of aluminium oxide.

The overall equation for the reaction is



- (a) Describe, in terms of electrons, the reduction and oxidation occurring during the electrolysis. [2 marks]

9 (b)(i) Aluminium ore contains 36% aluminium oxide, Al_2O_3 .

Calculate the mass of aluminium oxide in 500 tonnes of the ore. [1 mark]

Mass = _____ tonnes

9 (b)(ii)

Calculate the mass of aluminium metal that could be produced from this mass of aluminium oxide. [3 marks]



Mass = _____ tonnes

6

END OF PAPER


Question number	Additional page, if required. Write the question numbers in the left-hand margin.

Question number	Additional page, if required. Write the question numbers in the left-hand margin.

Question number	Additional page, if required. Write the question numbers in the left-hand margin.

Question number	Additional page, if required. Write the question numbers in the left-hand margin.

Question number	Additional page, if required. Write the question numbers in the left-hand margin.



GCSE

3410UB0-1

MONDAY, 22 MAY 2023 – MORNING

CHEMISTRY – Unit 2:

**Chemical Bonding, Application of Chemical Reactions
and Organic Chemistry**

HIGHER TIER

1 hour 45 minutes plus your additional time allowance

DIAGRAM BOOKLET

**This Diagram Booklet MUST be handed in to
the invigilators and sent for marking.**

Surname _____

First name(s) _____

Centre Number _____

Candidate Number 0 _____

DIAGRAM 1.1




TABLE 1.2

Part of the energy profile	Letter
energy change for the reaction	_____
energy of the reactants	_____
activation energy of the reaction	_____

DIAGRAM 2.1




TABLE 3.1



Name	Molecular formula	Structure	Homologous series
ethanol	C_2H_5OH		alcohols
ethanoic acid	CH_3COOH		
	C_3H_6		alkenes

DIAGRAM 3.2




DIAGRAM 3.3









DIAGRAM 3.4



THIS IS A BLANK PAGE
TURN OVER

TABLE 4.1

	Sodium chloride solution	Iron(II) chloride solution
zinc	no change	colour change
lead	no change	no change

TABLE 4.2

Mass of magnesium powder added (g)	Temperature (°C)
0	21.0
2.0	23.2
4.0	25.1
6.0	27.5
8.0	29.7
10.0	31.6

GRAPH 4.3

Temperature (°C)




TABLE 6.1

Solution	Reaction with magnesium
X	rapid fizzing, salt formed, temperature increase of 18 °C
Y	no reaction
Z	slow fizzing, salt formed, temperature increase of 11 °C

DIAGRAM 7.1




TABLE 8.1

Fruit juice drink	pH	Water (%)	Citric acid (%)	Ascorbic acid (mg/100 g)	Sugar (%)
lime	2.2	76.4	4.80	29	1.68
lemon	2.4	81.3	4.60	53	1.80
grapefruit	3.0	90.3	1.35	38	2.34
orange	3.6	87.4	0.96	50	4.85
tangerine	3.8	85.7	0.74	31	7.89



GCSE

3410UB0-1

MONDAY, 22 MAY 2023 – MORNING

CHEMISTRY – Unit 2:

**Chemical Bonding, Application of Chemical Reactions
and Organic Chemistry**

HIGHER TIER

Data Booklet

FORMULAE FOR SOME COMMON IONS

POSITIVE IONS	
Name	Formula
aluminium	Al^{3+}
ammonium	NH_4^+
barium	Ba^{2+}
calcium	Ca^{2+}
copper(II)	Cu^{2+}
hydrogen	H^+
iron(II)	Fe^{2+}
iron(III)	Fe^{3+}
lithium	Li^+
magnesium	Mg^{2+}
nickel	Ni^{2+}
potassium	K^+
silver	Ag^+
sodium	Na^+
zinc	Zn^{2+}

NEGATIVE IONS	
Name	Formula
bromide	Br^-
carbonate	CO_3^{2-}
chloride	Cl^-
fluoride	F^-
hydroxide	OH^-
iodide	I^-
nitrate	NO_3^-
oxide	O^{2-}
sulfate	SO_4^{2-}

THE PERIODIC TABLE

PERIODIC TABLE – KEY ATOMIC NUMBER – SYMBOL – NAME

1	H – Hydrogen
2	He – Helium
3	Li – Lithium
4	Be – Beryllium
5	B – Boron
6	C – Carbon
7	N – Nitrogen
8	O – Oxygen
9	F – Fluorine
10	Ne – Neon
11	Na – Sodium
12	Mg – Magnesium
13	Al – Aluminium
14	Si – Silicon
15	P – Phosphorus
16	S – Sulfur
17	Cl – Chlorine
18	Ar – Argon
19	K – Potassium
20	Ca – Calcium
21	Sc – Scandium
22	Ti – Titanium
23	V – Vanadium
24	Cr – Chromium
25	Mn – Manganese
26	Fe – Iron
27	Co – Cobalt
28	Ni – Nickel
29	Cu – Copper
30	Zn – Zinc
31	Ga – Gallium
32	Ge – Germanium
33	As – Arsenic

34	Se – Selenium
35	Br – Bromine
36	Kr – Krypton
37	Rb – Rubidium
38	Sr – Strontium
39	Y – Yttrium
40	Zr – Zirconium
41	Nb – Niobium
42	Mo – Molybdenum
43	Tc – Technetium
44	Ru – Ruthenium
45	Rh – Rhodium
46	Pd – Palladium
47	Ag – Silver
48	Cd – Cadmium
49	In – Indium
50	Sn – Tin
51	Sb – Antimony
52	Te – Tellurium
53	I – Iodine
54	Xe – Xenon
55	Cs – Caesium
56	Ba – Barium
57	La – Lanthanum
72	Hf – Hafnium
73	Ta – Tantalum
74	W – Tungsten
75	Re – Rhenium
76	Os – Osmium
77	Ir – Iridium
78	Pt – Platinum
79	Au – Gold
80	Hg – Mercury

81	Tl – Thallium
82	Pb – Lead
83	Bi – Bismuth
84	Po – Polonium
85	At – Astatine
86	Rn – Radon
87	Fr – Francium
88	Ra – Radium
89	Ac – Actinium