



GCE AS/A LEVEL

2410U20-1

CHEMISTRY – AS unit 2

Energy, Rate and Chemistry of Carbon Compounds

FRIDAY, 25 MAY 2018 – MORNING

1 hour 30 minutes plus your additional time allowance

Surname _____

Other Names _____

Centre Number _____

Candidate Number 2 _____

For Examiner's use only		
Question	Maximum Mark	Mark Awarded
Section A 1. to 7.	10	
Section B 8.	11	
9.	12	
10.	16	
11.	12	
12.	13	
13.	6	
Total	80	

ADDITIONAL MATERIALS

In addition to this examination paper, you will need a:

- calculator;
- DATA BOOKLET supplied by WJEC.

INSTRUCTIONS TO CANDIDATES

Use black ink, black ball-point pen or your usual method. Do not use gel pen or correction fluid.

Write your name, centre number and candidate number in the spaces on the front cover.

SECTION A Answer ALL questions in the spaces provided.

SECTION B Answer ALL questions in the spaces provided.

Candidates are advised to allocate their time appropriately between **SECTION A (10 MARKS)** and **SECTION B (70 MARKS)**.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

The maximum mark for this paper is 80.


Your answers must be relevant and must make full use of the information given to be awarded full marks for a question.

The assessment of the quality of extended response (QER) will take place in Q.10(a)(i).


If you run out of space, use the additional page(s) at the back of the booklet, taking care to number the question(s) correctly.

3. Explain why ethanol is soluble in water but ethane is not. [2]

4. Complete the diagram to show the formation of the π -bond in ethene. [1]



5(a) Complete the equation for the fermentation of glucose. [1]



(b) Name the substance generally used to catalyse this reaction. [1]

6. Draw TWO repeat units for the polymer formed from the monomer pent-2-ene. [1]

7. Draw diagrams to show the structures of the **E** and **Z** isomers of 2-bromopent-2-ene. Label the isomers **E** and **Z**. [2]

SECTION B

Answer ALL questions in the spaces provided.

- 8. A student was told that he could prepare chloroethane, C_2H_5Cl , by mixing ethane with chlorine. He added 2.0 g of ethane to excess chlorine and left the mixture exposed to ultraviolet light for several hours. He was then able to use a university laboratory to see whether chloroethane had been made.**
- (a) State an instrumental method by which the sample could be analysed. Explain how this would show that chlorination had occurred. [2]**

- 8(c) The student found that he had made 1.0 g of chloroethane. Calculate his percentage yield. [3]

Percentage yield = _____ %

8(d) The student was disappointed by the yield he obtained but his teacher told him that the yield was always poor due to other products being formed in this reaction.

Identify TWO other organic products, apart from chloroethane, and explain how they are formed.

[2]

9(a)(i)

The average bond enthalpy of a C—C bond is quoted as 348 kJ mol^{-1} .

Explain what is meant by BOND ENTHALPY. [2]

9(a)(ii)

Ethyne, C_2H_2 , contains a $\text{C}\equiv\text{C}$. It reacts with hydrogen in a similar way to ethene.




Some average bond enthalpies are given in the table.

Bond	Average bond enthalpy / kJ mol^{-1}
$\text{C}\equiv\text{C}$	839
$\text{C}-\text{C}$	348
$\text{C}-\text{H}$	413
$\text{H}-\text{H}$	436

Use the data to calculate the enthalpy change, ΔH , for the reaction of ethyne and hydrogen. [3]

$\Delta\text{H} =$ _____ kJ mol^{-1}



9(a)(iii)

Use your answer to part (ii) to sketch an energy profile diagram for this reaction on the axes opposite. Label ΔH and the activation energy, E_a , on your diagram. [2]

Substance	Enthalpy change of combustion, $\Delta_c H^\theta / \text{kJ mol}^{-1}$
hydrogen, H_2	-286
ethyne, C_2H_2	-1300
ethane, C_2H_6	-1600

9(b) Enthalpy changes of reaction are often found indirectly. The enthalpy change for the reaction of ethyne with hydrogen, as shown in part (a), can be determined by using enthalpy changes of combustion.

The table opposite gives some enthalpy changes of combustion, $\Delta_c H^\theta$.

Use these enthalpy changes to calculate the enthalpy change, ΔH , for the reaction of ethyne and hydrogen. [3]



$\Delta H =$ _____ kJ mol^{-1}

9(c) The theoretical values that you have calculated in parts (a)(ii) and (b) are both the enthalpy change for the reaction between ethyne and hydrogen.

Suggest a reason why these values are not the same. [1]

(d) Suggest the type of reaction that occurs between ethyne and hydrogen. [1]

12

10(a) Halogenoalkanes can be hydrolysed using water in a similar way to using aqueous sodium hydroxide.

A student carried out an experiment to investigate the rate of reaction for the hydrolysis of halogenoalkanes using water. The student used aqueous ethanol to dissolve the halogenoalkane and then added a few drops of aqueous silver nitrate. He timed how long it took to produce a precipitate. He obtained the results shown in the table.

Halogenoalkane	Time / s
1-chloropropane, C₃H₇Cl	300
1-bromopropane, C₃H₇Br	90
1-iodopropane, C₃H₇I	15

Bond	Bond enthalpy / kJ mol⁻¹
C—H	413
C—C	348
C—F	485
C—Cl	328
C—Br	276
C—I	240

Element	Electronegativity
chlorine	3.16
bromine	2.96
iodine	2.66
carbon	2.55

10(a)(ii)

Write an IONIC equation, including state symbols, for a reaction that produces a silver halide precipitate. [1]

(iii) Suggest a practical method by which the student could have obtained these results. [2]


(iv) Suggest the difference that the student would have observed in his experiments if he had not added ethanol to the water before adding the halogenoalkane. [1]

10(b)(ii)

Use relevant data in part (a) to explain why hydrofluorocarbons, HFCs, have replaced CFCs in many of their uses. [2]

16

Intensity




m/z

11. Compound **X** contains only carbon, hydrogen and oxygen. On analysis it was found to contain 53.3% carbon and 11.1% hydrogen by mass.

A simplified form of the mass spectrum is shown opposite. The infrared absorption spectrum for **X** is shown opposite page 25.

The low resolution ^1H NMR spectrum of **X** has three peaks.

When **X** is warmed with excess acidified potassium dichromate(VI) there is a colour change. The organic product of this reaction does NOT react with aqueous sodium carbonate.



11(b) (i) State the type of reaction that occurs when **X** is warmed with acidified potassium dichromate(VI). [1]

(ii) Draw the structure of the organic product formed when **X** reacts with acidified potassium dichromate(VI). [1]

12(a) Iodide ions can be oxidised to iodine by reaction with acidified hydrogen peroxide.



The rate of reaction can be followed in a clock reaction by the appearance of a blue-black colour.


An experiment was carried out to determine the effect on the rate of reaction of varying the concentration of iodide ions. All other volumes and concentrations were kept constant. The results are shown in the table on page 29.

12.

Concentration I^- / mol dm ⁻³	Time for appearance of blue-black colour / s	Rate / s ⁻¹ × 1000
0.1	56	18
0.2	20	_____
0.3	18	_____
0.4	12	_____
0.5	10	_____

12(a)(i) Use rate = $\frac{1000}{\text{time}}$ to calculate the rate for each experiment and complete the table above. [1]

(ii) On the axes opposite, plot the concentration of I^- against rate and draw a suitable line. [3]



12(a)(iii)

From the graph deduce the relationship between concentration of I^- and rate of reaction. [1]

(iv) **USE THE GRAPH** to calculate the time the reaction would take to turn blue-black using a 0.15 mol dm^{-3} solution of I^- . Show clearly how you obtained your answer. [2]

Time = _____ s

12(a)(v)

For each experiment the rate was calculated using the time taken to produce excess iodine. Explain why this is only an approximation for the rate as the reaction proceeds. [2]

(b)(i) Draw a Boltzmann energy distribution curve. Label the axes. [2]


12(b)(ii) Use this energy distribution curve to explain how catalysts affect the rate of a reaction. [2]

13

13(a) Explain what is meant by a carbon-neutral fuel. [3]

13(b) Ethane and an unknown alkane were burned in oxygen.

It follows from the equation below that one volume of ethane produced two volumes of carbon dioxide and three volumes of water vapour.



10 cm³ of the unknown alkane C_xH_y burned according to the following equation.



The total volume of carbon dioxide and water vapour produced was 20 cm³ more than the original volume of C_xH_y and oxygen. All volumes were measured at the same temperature and pressure.

13(a)(i)

State the volumes of carbon dioxide and water vapour produced on burning 10 cm^3 of the unknown alkane in terms of x and y . [1]

Volume carbon dioxide = _____ cm^3

Volume water vapour = _____ cm^3

(ii) Calculate the value of y . [2]

$y =$ _____

6

END OF PAPER

